

III. have been subjected to repeated tests by me in the venereal wards of the Charity Hospital, as well as in private practice.

In my studies, clinical and experimental, I have received valuable assistance from my friend Dr. L. A. Stimson, and from Dr. Dinslow, of the House Staff of the Charity Hospital.

E. L. KEYES.

210 MADISON AVENUE, NEW YORK, *January*, 1877.

CONTENTS.

CHAPTER I.

GENERAL TREATMENT OF SYPHILIS.

A Rational Treatment of Syphilis requires Stronger Support than Simple Statistics.—All Cases, light as well as severe, require Continued General Treatment, since Cases commencing mildly may become severe, and since the General Treatment proposed is tonic to the Individual.—Proof of the Tonic Action of Small Doses of Mercury: *rational*, from Clinical Experience and the General Sanction by the Profession, of the Use of Minute Doses of Mercury as an Alternative in Chronic Disease; *scientific*, from Actual Blood counting with the Hématimètre and Experiments upon Animals.—Description of the Hématimètre, the Method of using it, Fluid to be employed, etc.—Effect of Mercury and of the Iodides upon the Blood as shown by the Hématimètre.—Proof that the Mercury controls the Symptoms of Syphilis . . . 7

CHAPTER II.

DETAILS OF GENERAL TREATMENT.

When to commence General Treatment.—Details of the Continuous Tonic Treatment by Mercury.—The Tonic Dose; the Full Dose; the Reserve Dose.—The Duration of General Treatment.—Inunction; Subcutaneous Injection; Mercurial Vapor-Bath.—Treatment of Salivation.—Preparations of Iodine and their Use in Syphilis.—Mixed Treatment.—Limit to the Dose of the Iodides.—Duration of Treatment by Iodine.—Other Means used in General Treatment: the Woods, Zittman's Decoc-tion, Syphilization, Hot Springs of Arkansas.—Treatment of Pregnant Women.—Treatment of Inherited Disease 31

CHAPTER III.

THE LOCAL TREATMENT OF SYPHILIS, AND THE ESPECIAL MEANS
ADAPTED TO SPECIAL LESIONS.

Local Treatment of Chancre.—Of Syphilitic Phagedena, of Bubo.—Of
Lesions of Mucous Membranes.—Of Cutaneous Lesions.—Of Lesions
of the Eye, the Ear, of Muscles, Joints, Bones.—Visceral and Nervous
Syphilis 66

SYPHILIS.

CHAPTER I.

GENERAL TREATMENT OF SYPHILIS.

A Rational Treatment of Syphilis requires Stronger Support than Simple Statistics.—All Cases, light as well as severe, require Continued General Treatment, since Cases commencing mildly may become severe, and since the General Treatment proposed is tonic to the Individual.—Proof of the Tonic Action of Small Doses of Mercury: *rational*, from Clinical Experience and the General Sanction by the Profession, of the Use of Minute Doses of Mercury as an Alternative in Chronic Disease; *scientific*, from Actual Blood counting with the Hématimètre and Experiments upon Animals.—Description of the Hématimètre, the Method of using it, Fluid to be employed, etc.—Effect of Mercury and of the Iodides upon the Blood as shown by the Hématimètre.—Proof that the Mercury controls the Symptoms of Syphilis.

A RATIONAL treatment of syphilis must rest upon a surer foundation than the mere statement of its value made by him who employs it. Such a statement has been made again and again, by different people employing the most varied means against the same evil; and, with apparent justification, for it is well known that a majority of the most visible lesions of syphilis (the cutaneous efflorescences) tend to subside spontaneously, and thus, as Fournier happily expresses it, to afford a triumph to every mode of treatment.

There still exist ardent advocates of syphilization,