

## LECTURE XXIII.

- NERVOUS COMPLICATIONS OF FEVER, *continued*—Cerebro-spinal fever—Phenomena of fever inconstant and variable, except, perhaps, the phenomenon of increased temperature—Type of fever also varies in different epidemics—Two examples: (1) yellow fever of 1826-27, (2) malignant purpuric, or cerebro-spinal fever of 1867—Dr. E. W. Collins' report on latter—There exists a "constitutional element" in the disease, so that the cerebro-spinal arachnitis can hardly be held to be a primary, idiopathic affection—Evidences of essentiality from presence of other phenomena in connection with the skin, etc.—Reports to the *Medical Society of the King and Queen's College of Physicians in Ireland* on the epidemic of 1867—Inconstancy and variability of the symptoms in the outbreak—Dr. H. Kennedy's views—Symptoms of the disease—Petechiæ—Early setting in of putrefaction—Retraction of head; sometimes persistent after disappearance of other local and general symptoms, and sometimes persistent after death—Recapitulation: Points to be considered in connection with epidemic of 1867: (1) yellow fever of 1826-27, (2) cerebro-spinal arachnitis of 1846 (Dr. Mayne), (3) coincidence of cases of malignant measles in 1867, and (4) hemorrhagic and purpuric smallpox in epidemic of 1871-72 . . . . . 165

## LECTURE XXIV.

- NERVOUS COMPLICATIONS OF FEVER, *continued*—Hysteria—Occurrence of hysteria, especially at an early stage, of unfavourable import—View that hysteria is always symptomatic of uterine excitement is quite erroneous—Nymphomania only a local and accidental manifestation—Hysteria is observed in males as well as in females in fever—Case of erotic symptoms in typhoid fever occurring in a young girl, reported by Dr. A. W. Foot—In early stage of fever hysteria generally is the precursor of severe nervous symptoms—Its appearance may lead to serious complications later in the disease—Illustrative cases—Hysterical symptoms are sometimes connected with actual or organic disease, especially in acute affections—Dr. Cheyne's observation: *Hysteria a ground for a good prognosis in every disease, fever alone excepted*—Outbreak of hysteria, affecting the abdomen, in female fever ward of Meath Hospital—Anomalous symptoms in advanced stages of fever often due to hysterical state—Case of typhoid hysteria in the male followed by cerebritis . . . . . 173

## PART II.

## TREATMENT OF FEVER.

## LECTURE XXV.

- INTRODUCTORY REMARKS—Principles on which the treatment of fever is to be based—True meaning of the word *empiric*. Historical retrospect—The Symptomological, the Anatomical, the Rational or Eclectic Schools—Essence of fever cannot be determined by pathological anatomy—*Etiology* of fever is indefinite . . . . . 177

## LECTURE XXVI.

- No specific line of treatment—Respect to be had (1) to the essential disease, (2) to its local and secondary effects—Failure of specifics in early stage of fevers—Want of success in the endeavour to found a science of therapeutics on experimental physiology or pathology—Effects of the action of the law of periodicity wrongly attributed to the adoption of therapeutical measures—Sustenance by food and stimulants—Two sources of danger to the fever patient: (1) primary effects of the fever poison in causing depression, (2) supervention of secondary local disease—Views of Dr. Graves on the subject of giving food in fever . . . . . 185

## LECTURE XXVII.

- STIMULANTS IN FEVER—Views as to the nutrient properties of stimulants are to be received with caution—*Anticipative* use of stimulants—Meaning of the term—Considerations to be taken into account in resolving upon this method of treatment: (1) prevailing epidemic character of the disease, (2) previous condition of the patient—"Sinking of vital power"—Illustrative case—Stimulation often unsuccessful in the intemperate, and in those whose brains are over-worked, (3) development of symptoms of severe typhus, (4) development of fever odour—Contrast between typhus and typhoid as regards period at which stimulation is called for—*Condition of the heart, a guide*—Physical signs of cardiac weakening . . . . . 193

## LECTURE XXVIII.

- STIMULANTS IN FEVER, *continued*—Signs in connection with the heart of the agreement of stimulants: (1) return of impulse, (2) return of first sound, (3) gradual fall in the rate of the pulse—In cases of "fœtal heart" great boldness in stimulation is needed—No certain rules as to quantity of wine and whiskey or brandy required—Examples of free use of stimulants in malignant typhus—Case of Harcastle (typhoid fever)—Eruption of vesicles as a secondary complication—Bed-sores . . . . . 201

## LECTURE XXIX.

- STIMULANTS IN FEVER, *continued*—Case of Harcastle, *continued*—Treatment by food and stimulants in extreme cases—Presence of cerebral symptoms to a great extent unfavourable to the exhibition of stimulants—Necessity for daily observation to the effects of the treatment in each case—Signs of disagreement of stimulants—*Routine* practice is in every instance to be deprecated—Fallacies of the numerical system in therapeutics—History of *routinism*—Its results—Description of *routinism* in the treatment of fever . . . . . 212

## LECTURE XXX.

- TREATMENT OF THE LOCAL SECONDARY AFFECTIONS IN FEVER—Relative importance of these affections as regards *prognosis*—BRONCHIAL AFFECTION—Necessity for administration of stimulants and nourishment—Danger of exhibition of tartar emetic—Failure of emetics—Turpentine-punch—Dry-cupping, poulticing, blistering—*Internal remedies*: bark, ammonia, spirit of chloroform, turpentine—ACUTE CONSOLIDATION OF THE LUNG—Its three forms—Treatment of the first form by dry-cupping, blisters, quinine, turpentine, and wine—Of the second form by local depletion simultaneously with the administration of wine—Of the third form, *externally* by iodine and blisters, *internally* by tonics and iodide of potassium . . . . . 222

## LECTURE XXXI.

- TREATMENT OF INTESTINAL SECONDARY AFFECTIONS—Two chief indications: (1) alleviation of symptoms, (2) modification of typhoid deposition—Poulticing—Local depletion in early stage—Analogy in variolous eruption—Danger of alterative or purgative treatment at the outset of Continued Fever—Necessity for caution—*Constipation*—*Diarrhœa*—Poultices, demulcents, sedative astringents, injections of flax-seed tea—*Tympany*—Turpentine injection—*Diet* in diarrhœa—*Perforative peritonitis*—Opium our sheet-anchor—Danger of the antiphlogistic method—Dr. Murchison on the treatment of this accident—Bran poultices and warm fomentations—*Hemorrhage* from the intestine in fever—Not to be interfered with unless continued and excessive—Treatment by astringents, opium—Illustrative case . . . . . 227

## LECTURE XXXII.

	PAGE
TREATMENT OF THE NERVOUS SECONDARY SYMPTOMS OF FEVER—HEADACHE—Cold lotions, warm fomentations, moderate leeching, shaving the head, cold affusion, ice—DELIRIUM—Treatment depends on (1) period of case, (2) presence of hyperæmia of the brain, or otherwise—Ice, leeches, shaving the head, cold affusion in <i>active</i> delirium—Nourishment and wine in <i>passive</i> or anæmic delirium—SLEEPLESSNESS—Moderate leeching, cold affusion, ice—Turpentine in constipation and tympany—Catheterism in distended bladder—Sedatives—Opium, tartar emetic and opium, hyoscyamus, bromide of potassium, chloral, wine—CONVULSIONS—Most formidable in fever—Uræmic, due to (1) <i>retention</i> of urine: catheterism; (2) <i>suppression</i> of urine: dry cupping and poulticing over kidneys, diluents, diuretics, aperient enemata, promotion of action of the skin	237

## LECTURE XXXIII.

PHLEGMASIA ALBA DOLENS—The swelling is not always painful, or white in appearance—Symmetry of the affected limb not lost—Professor Trousseau's views as to the etiology of the affection—Phlegmasia (1) of puerperal women, (2) in scrofulous and (3) cancerous cachexia—Pulmonary <i>embolism</i> caused by phlegmasia—Case of phlegmasia after typhus fever—TREATMENT of the affection—GENERAL CONCLUSION	245
APPENDIX B	253
INDEX	257

## LECTURES

ON

THE DISEASES OF  
THE NERVOUS SYSTEM.

DELIVERED AT LA SALPÊTRIÈRE

BY

J. M. CHARCOT,


PROFESSOR TO THE FACULTY OF MEDICINE OF PARIS; PHYSICIAN TO LA SALPÊTRIÈRE; MEMBER OF THE ACADEMY OF MEDICINE, AND OF THE CLINICAL SOCIETY OF LONDON; PRESIDENT OF THE ANATOMICAL SOCIETY, AND EX-VICE-PRESIDENT OF THE BIOLOGICAL SOCIETY OF PARIS, ETC.

TRANSLATED FROM THE SECOND EDITION BY

GEORGE SIGERSON, M.D., M.Ch.,

LICENTIATE OF THE KING AND QUEEN'S COLLEGE OF PHYSICIANS; LECTURER ON BIOLOGY AND EX-DEAN OF THE FACULTY OF SCIENCE, CATHOLIC UNIVERSITY OF IRELAND; FELLOW OF THE LINNEAN SOCIETY OF LONDON; MEMBER OF THE SCIENTIFIC SOCIETY OF BELGIUM, AND OF THE ROYAL IRISH ACADEMY, ETC.

WITH ILLUSTRATIONS.



PHILADELPHIA:  
HENRY C. LEA.  
1879.