

more difficult to determine than in the male, for desire is more variable and excess as a rule is better borne. There should certainly be a respite, however, during menstruation, in the latter period of utero-gestation, and for several weeks after confinement.

Copulation may be incomplete from various causes; it is not infrequently so in the first experience of either party to it. The undue nervous excitement of the male sometimes ends in a loss of rigidity or in premature ejaculation. An unduly rigid hymen, an undeveloped condition of the vagina, narrowness of the orifice, or other abnormality may prevent entrance.

Various accidents occur. Lacerations and abrasions of the penis are usually of no importance except with reference to the possibility of infection. Edema or ecchymosis of the skin occasionally follows undue violence. Urethral hemorrhage sometimes occurs, frequently from rupture of a stricture. Cases are reported in which the root of the penis has been dislocated. The so-called fracture consists in a rupture of the corpora cavernosa and may result in such deformity as to render future coitus impossible. Quite a number of cases have been recorded in which the great contraction of the vagina aided by that of the levator ani rendered separation after coitus difficult or impossible until various remedies had been applied. This result seems to have followed the application of musk to the vagina in several instances. More or less complete epileptic seizures sometimes occur during or after connection, most frequently accompanying the orgasm, particularly in the female. Death has also been reported in not a few instances, but its immediate cause has generally been cerebral hemorrhage, the rupture of an aneurism, or it was the termination of an organic disease hastened by the exertion.

The injuries to the female are sometimes serious in character; they are especially liable to occur as a result of the assumption of unnatural positions. The rupture of the hymen is generally of no consequence, but the rent occasionally extends so deeply as to involve the mucous membrane of the vagina and thus give rise to serious hemorrhage. Laceration of the perineum, the clitoris, the urethra, or the anterior vaginal wall into the bladder and similar accidents have been reported. Uterine hemorrhage is not seldom induced, especially by the first connection when it occurs soon after menstruation, and is sometimes of considerable severity.

The medico-legal aspects of this subject will be found under the title of *Rape, Medico-Legal Aspects of*.  
James M. French.

**COPYRIGHT.**—The copyright law of the United States does not accord any peculiar privileges to authors of medical works. Such works may be copyrighted like any others. The chief requirements are: 1. Before publication mail a printed copy of the title to the Librarian of Congress, Washington, D. C. 2. Upon the title page or its back, print the customary notice that the book is copyright. 3. Within ten days after publication send two copies, by mail or otherwise, to the Librarian. A letter of inquiry addressed to the Librarian will bring a printed circular giving full directions as to details. Compliance with the prescribed conditions secures, if the author is a citizen or resident of the United States, the sole liberty of publishing the work for twenty-eight years; and there is a qualified right of renewal which, in the case of a work of permanent character, is valuable. A book duly copyrighted cannot be republished in full, or copied in its substantial or important parts. But it is not every use of a treatise, on the part of subsequent writers, that is considered an unlawful infringement of its copyright. Limited and reasonable use of the contents of earlier works, though copyrighted, by way of study or quotation of extracts, is allowable. Moreover, there is a clear distinction between a book as a literary property and the science which it explains. Copyrighting a text-book on the composition and use of medicines does not secure an exclusive right to compound the medicines described, not even if they are newly devised by

the author; the very object of publishing a medical work is to communicate to practitioners the knowledge of symptoms and prescriptions which it contains, for their use in current practice. Photographs skillfully showing diseases or deformities, etc., may be copyrighted. Mere labels, such as are placed upon bottles to designate the medicines which they contain, and the diseases for which they are adapted, are not within the protection of the copyright law; such labels often bear notice that they are "copyright," but this is nugatory. Medical lectures, if unpublished, are not, strictly, a subject of copyright; for the required two copies cannot be supplied to the Librarian of Congress. But they are protected by a statute which forbids publishing any manuscript without obtaining the consent of the author or proprietor. Oral delivery of a lecture to a class of students does not give them the right to take notes for any purpose of publication, or of selling copies to other persons; the right of members of the class is limited to memorizing and taking notes of the lecture for their individual instruction and guidance in future practice. The copyright law important to medical writers and publishers is embodied in "United States Revised Statutes," Title IX., Chapter 3. Drone on "Copyright" is the most recent law treatise. Morgan's "Law of Literature" is comprehensive and readable. Abbott's "National Digest," title "Copyright" states the decisions and statutes to 1884, with a legal bibliography.  
Benjamin Vaughan Abbott.

**CORAL BEAN.** See *Poisonous Plants*.

**CORIANDER.**—CORIANDRUM. Coriander Fruit; incorrectly, "Coriander Seed." The coriander plant is an annual European herb, also cultivated in Europe and the United States. It has been so long cultivated that it is



scarcely known in an indubitably wild state. The following is the official description of the drug: Globular; about 4 mm. in diameter; crowned with the calyx-teeth and stylopod; brownish yellow, with slight longitudinal ridges; the



FIG. 1515.—Coriander Plant, one-tenth natural size. (Ballion.) FIG. 1516.—Fruit of Coriander, natural size.

two mericarps cohering, enclosing a lenticular cavity, and each furnished on the face with two oil-tubes; odor and taste agreeably aromatic.

It contains about one-half per cent. of essential oil, eleven of fat (in the seeds), and fourteen of nitrogenous matters, etc. The only constituent of value is the oil (*Oleum Coriandri*, U. S. P.), a colorless or yellowish liquid, having the characteristic aromatic odor of coriander, a warm, spicy taste, and a neutral reaction, sp. gr. 0.870 to 0.885. *Linatool*, or *coriandrol*, and *pinene* are the constituents.

Coriander is a very mild and agreeable carminative, of the same properties as anise. Dose 2 to 8 gm. (3 ss.-ij.); of the oil, ten drops or so—on sugar or suspended in syrup or mucilage. There is no official preparation of either the fruit or the oil, though the former enters into the *Confectio sennæ* and the latter into the *Compound Spirit of Orange*.  
W. P. Bolles.

**CORIARIA.**—A genus (fam. *Coriariaceæ*) of about ten species of shrubs inhabiting warm or tropical regions. The leaves and fruits of several of these are well-known poisons, both to man and to grazing animals. In the ripe fruit, the poison is said to be confined to the seeds. Their medicinal use has apparently not been tried, but they are of interest because the leaves of one, *C. myrtifolia* L., have been found as an adulterant of senna. They are readily distinguished by their three-nerved surface. The active principle is the glucoside *Coriomyrtin* ( $C_{20}H_{30}O_{10}$ ), which is crystalline, tasteless, and soluble in both water and alcohol. Its action is very similar to that of picrotoxin.  
Henry H. Rusby.

**CORNEA, DISEASES OF THE.**—In order to recognize and understand the various pathological lesions to which the cornea is subject, an accurate knowledge of its anatomy and histology is essential. The term *keratitis* is applicable to all inflammatory processes affecting the cornea and causing certain deviations from its normal uniform transparency,—alterations which are, as a rule, easily recognized by simple inspection with the naked eye; but the method of examination known as focal illumination affords a ready means of detecting changes of slight degree which might otherwise easily escape observation. Structural changes resulting from inflammation are either transitory or permanent. Inflammatory processes in the cornea give rise to various changes in its physical condition, the most constant of which is a loss of transparency which involves the compact fibrillar arrangement of the corneal tissue, and which is caused by the inhibition of fluid material as well as of the more solid cellular elements that have migrated into the cornea from the surrounding vascular system. This subject has been very thoroughly studied by many observers, and the free migration of leucocytes into the cornea from this source is one of the pathological questions which may be regarded as definitely settled. The formation of blood-vessels in the cornea is common to several varieties of keratitis. They owe their origin to the fact that the terminal vessels surrounding the cornea send off-shoots into the corneal tissue and so give rise to a conspicuously vascular condition of this structure, which disappears wholly or in part as the inflammation subsides. The development of these blood-vessels, therefore, belongs to the process of repair. An engorgement of the blood-vessels in the tissues around the cornea is one of the most constant signs of keratitis, and is known as pericorneal congestion.

It is still an open question as to what part the fixed corneal corpuscles take in the inflammatory process. If this has been of such a character as to cause permanent alteration in any part of the cornea, both the fixed corpuscles and the fibrillar structure involved are found to have undergone more or less destruction and elimination. A transformation of the normal clear corneal tissue into white fibrous or scar tissue, which often remains permanently, is a common sequel of corneal inflammation; this is the explanation of most permanent corneal opacities.

The cornea, owing to its exposed position and non-vascularity, is particularly liable to traumatism and to inflammatory processes due to infection and to lessened vitality of the tissues. Three types of corneal inflammation are generally recognized; they are *infiltration*, *abscess*, and *ulceration*.

1. *Infiltrations* are characterized by the presence of leucocytes in such numbers as to cause loss of transparency, with more or less displacement of the fibrous parenchyma of the cornea. Crowding the lacunæ and canaliculi, they may give rise to an ill-defined, streaky opacity, or, when they are accumulated in larger masses, the cloudiness may be more uniform and circumscribed. Notwithstanding this the corneal tissue may remain so far intact as to permit of perfect restoration, when resolution takes place. If, however, the process has been of long duration and considerable intensity, some degree of permanent opacity will result; the migratory cells may indeed become transformed into fibrous tissue (sclerosed),

and blood-vessels may become developed in the substance of the cornea as a part of the permanent alteration of its structure. Corneal infiltrations may be superficial or deep-seated, circumscribed or diffuse; and, although the tendency is to recovery without destruction of tissue or loss of transparency, the more unfavorable terminations in sclerosis or ulceration are quite common events, especially if the infiltration has been injudiciously treated with remedies which exaggerate the irritation already existing. Superficial infiltrations of the cornea are apt to cause more irritation than deep-seated ones, on account of the corneal nerve filaments being most abundant toward its anterior surface. This explains why we often see most intense irritation following trivial abrasions of the surface.

2. *Abscess of the cornea* occurs when an infiltration reaches such a degree of intensity that the nutrition of the part is interfered with to the extent of destroying the proper corneal tissue, while the infiltrating elements become transformed into pus cells. The distinctive gross characters of an abscess are its circumscribed appearance and yellowish color. It often gives rise to an accumulation of pus in the anterior chamber (*hypopyon*), presumably in consequence of the fact that the irritant toxins evolved in the inflamed area find their way into the anterior chamber by absorption or transudation (just, for instance, as atropine solution does when dropped into the conjunctival sac), and there excites a purulent iritis or descemetitis. The proof of this lies in the fact that although pyogenic bacteria are as a rule found in abundance in the purulent corneal lesion, there are few if any of these in the accompanying hypopyon. So much of the corneal tissue as may have been actually destroyed can only be imperfectly regenerated; for which reason, more or less of an indelible opacity remains. Then again, as already stated, an abscess near the anterior surface of the cornea is more painful than one that is deep-seated. The duration of corneal abscess is variable; sooner or later the surface is apt to give way, and the abscess becomes transformed into an ulcer. More rarely it breaks through into the anterior chamber.

3. *Ulcers of the cornea*, many varieties of which have been described, all have one feature in common, that is, they are accompanied by loss of substance. They may result from an infiltration or an abscess, or the ulceration may be the primary lesion, commencing as a superficial loss of substance of irregular outline, first of the epithelium, then speedily extending through Bowman's membrane to the cornea proper. Deposits of lymph and pus in the anterior chamber are also quite common in certain types of corneal ulcers. The surface of the ulcer, in its progressive condition, is generally covered with an opaque material, the residue of broken-down tissues, while the margins are surrounded with a grayish zone of infiltration. When the healing process sets in, the base of the ulcer becomes cleaner, its margins smooth and rounded, and the peripheral opaque zone less conspicuous. As the healing progresses the ulcer becomes shallower, and its surface smooth and glistening; for at this stage it is partly covered with epithelium which has grown from the margins toward the centre. Beneath this, new tissue is developed, and the excavation due to loss of substance is gradually filled up by the process of cicatrization. There remains, finally, an opaque area of variable depth and extent, from a slight cloudiness (nebula) to a dense white cicatrix of a pearly or tendinous character (leucoma). Corneal ulcers, especially when situated near the margin, frequently become vascularized during the healing process. An ulcer may remain stationary for a long time, or the surface may not attain the level of the surrounding cornea (corneal facet). Sometimes the process of repair is arrested and renewed ulceration takes place. Perforation of the cornea, with adhesion or prolapse of iris, is a common occurrence, in which case the healing process is apt to be followed by yielding of the softer reparative material to the intraocular pressure and the development of a protrusion known as a corneal staphyloma. Extrusion of the crystalline lens is to be feared