



IOANNES TERRENTIVS
MAGISTER ET MERCHANTIAE
MEXICAE MERTORIQVE

HABES & hic, curiose Lector, quibus oculos satrem, si mi-
nus animum satiare valeas, plantas inquam à nostri orbis
flore præcipue & fructu differentes stirpibus, adeo multa &
mira adhuc in arario suo abscondita claudit diues Natura,
quæ Indies Europæis nostris cum voluptate ostendit. Fructu
interca nuda plantarum harum figura & partium delineatio-
ne, & colorum descriptione, quam ego ex Autographo dili-
genter excerpti. Viræ ab Auctore non additæ fuerunt, quod forsân adhuc Mexica-
no populo nondum cognita essent. Non omnia simul & semel Natura profudit.
Multa nunc patent quæ olim latebant, & harum non dubitandum quin stirpium ali-
quando vera experimenta habituri simus.



ALIA

ALIA RVM
NOVAE HISPANIAE
PLANTARVM
NARDI ANTONII RECCHI
IMAGINES, ET NOMINA.

ACAXAXA.



ACAXOCHITL.



IOANNIS TERRENTII LYNCEI NOTÆ.

ACAXAXA.

Ex singulis geniculis inferioribus
fibras in solum demittit folia
Gentianæ similia in nodis sin-
gulis totum caulem ambiunt in
summitate ex tribus folijs exit
iulus longus, colore ex viridi fla-
uescente.

ACAXOCHITL.

Limonia similis, vel Echio; caulis
ruber, folia longa Bistorta figu-
ra, flores rubei calycis foliola
longa, flos barbâ habet lon-
gam trium foliorum.

ACXO-



ACXOYATL.

Radix coloris castanei, ramuli sanguinei, folia minutissima alata in ramis, Pino similia.

AHATE DE PANVCHO:

Vel Ate Pannicensis, aut Tzypipatlis fructu apud Mexicanos præcellenti arbor nobilis. Folium Aurei Pomi, fructus cæruleus, sed in nigrum tendens, ferro polico similis, rotundus totus quasi ex glandibus constans, Pinæ nûcis figura.

Ramus vero coloris castanei, folia inglandis.


AMACPATLI

Coralli species, rami sunt argillæ colore, puncta e nigro rubent, filamenta omnia rubra.

ACXO



AMACPATLI.



NOTATIONES.
 AMOZOTL.



ACATZAN AYCXITL.


Similis Equifeto, radix meris capillis ex flauo nigrantibus constat. caules vero amethystini coloris.

AMAZAVHTLI.

Vel Amutzauhli, bulbacea è radice, folia crassa, Tulipis similia.

AMOZOTL.

Folia plerumque bina, Crithmi, longa, digiti latitudine, in summitate uua. Ligustro similes bacca, quarum color viridis ad obscuritatem tendens.



IOANNIS TERRENTII LYNCEI

AMAZQVILITL.

ANONYMA MECHV. II.

Videtur esse arbor. stipes pullus. rami habent folia Phyllireæ latifolia, unde oriuntur bacca virides, breui annexæ pediculo.

ANONYMA MECHVAC I.

Radix constat multis fibris nigris, caules lutei, folia latitudine sua adnascuntur. serrata sunt, vt Sideritis ferme. in summitate quid viride triangulare conspicitur.

ANONYMA MECHV. II.

Rami habent folia duo, parua, vna cum magno in quouis exortu. flores in summitate oblongi, quasi in umbella compositi sunt.

ANO-



NOTATIONES.

ANONYMA VI.

ANONYMA MICHVACANENSIS III.

Surculos multos virentes emittit, folia Menthae quadamtenus amula, ex atro virefcentia, serrata. Luteos flosculos oris purpurascens. Calyces virides. Radicem habet fibrosam, ex flauo ad fuscum vergentem.

ANONYMA IV.

Hæc in pictura, longa, & crassa radice conspicua est, ex sulco flauo rubente. in qua extuberant partes nonnullæ, & à lateribus fibræ demittuntur, folia quasi Coluteæ.

ANONYMA V.

Radix oblonga est crassitudinis, spadiceo coloris. folia saturate virent, venis albicanibus. flores fere purpurei.

ANONYMA VI.

In ramulo omnia oblonga. folia, flores, pediculi, calyculi florum suave rubent. ora lutea, exertis quasi linguis. folia fere Serratulae ex saturato virent.

Gg 2 ANO-



IOANNIS TERRENTII LYNCEI

ANONYMA VII.

Bulbosa videtur, fibroso è capite subflavo. folia breviora, sed Gladioli ferè. cauliculum promit geniculatum, longum, purpureum. bini ad nodos flores purpurei pariter pendent.

ANONYMA VIII.

Ex Maluaco genere Anglicanum Bami æmulari videtur colore, & venis. folia respondent, magis tamen cordata. Rubent flores, & ad oras dilutius ad calycem albicant.

ANONYMA IX.

Ramus tantum vt in superiore pingitur, concolor, namque flores & folia pariter virent.



ANONYMA IX.



ANO-



NOTATIONES.

AOCOXOCHITL.



ANONYMA X.

Radix crassa, & rubra, in flauum vergens, caulis ruber, folia bina, eminent flores in summo, similes Plumbagini.

APANXALOA. TA

Folia bina in quouis exortu foliorum apposita, Saponaria: similia. Caules rubri. flores longi, rubri aliquantum, carulei Vaccaria: similes.

AOCOXOCHITL.

Serpit hæc herba, & ex quouis nodo emitit fibras in terram. folia bina Vaccaria: similia. exoritur quid rubrum ex quouis ala, sed quid sit, non conspicitur.

Gg 3 AOVIT-

AQVITZTLI.



ATEHVAPATLI.



IOANNIS TERRENTII LYNCEI

AQVITZTLI.

Scandit hæc herba, Caule rubro, maculis nigris conserpo. quibus ramus tribus constat foliolis, instar Clematidis. folia etiam obscure rubent.

ATEHVAPATLI.

Radix rotunda est Aristolochia instar, folia bina, figuræ ovalis, nervi tantum tres, in longitudinem folij excurrentes: reliqui verò in transversum diuicata, ex quavis ala folij quid enascitur.

ATEXCALQVAHOITL.

Rami quasi lignei, coloris obscuri. Suberis adnata sunt, excrementa illa marina alba, punctis rubris, nigrisve variè conserpta.

-TIVOA

ATEN-



ATENXIHVITL.

355
ATZOYATL COZTIC YNI RODERILYO.

MONOTATIONES.

AXIXPATLI.

ATENXIHVITL, seu HENXIHVITL.

Trifolia humilior planta, ferratis folijs, fibrosa radice ex fusco pallida. pediculi foliorum rubent dilutius.

ATZOYATL COZTIC YNI RODERILYO.

Crassa, magna, magnisq. fibris radix, & spadicei coloris. folia longa, & ferrata. flores Conyzae lutei.

AXIXPATLI.

Folia bina, quamvis sculptor aliquantulum in hoc defecerit. Flores lutei, calyculis virentibus. Radix dilutius rubet.

-AYA

ATZL-



ATZIZICAZTLI.

AXIHVITLAPAPATLACTLI.



IOANNIS TERRENTII LYNCEI

AXOCHIOTL.

ATZIZICAZTLI.

Folia Vrticæ non bina, rami omnes raris
habent folia, Vrticæ similia; & ex
quavis ala enascuntur alia.

AXIHVITLAPAPATLACTLI.

Folia bina ex aduerso posita, pediculo
satis longo harentia. foliorum tum
longitudo tum, latitudo sex digito-
rum est.

IITATXIXA
AXOCHIOTL.

Hæc herba videtur serpere, rami folia
cordis figura, & repanda habent. si-
tu liguminum, siue alato, ita vt in
extima tria sint folia.

IXTA

AYA-



AYACACHQVAHOITL.

AZTAXOCHITL.



NOTATIONES.

BANAICXITL.

AYACACHQVAHOITL.

Folia Gossipio similia, sublutea, in sum-
mo multa simul congesta.

AZTAXOCHITL.

Quiuis ramus habet tria folia Vrticæ si-
milia alte incisa ordinata vt in Pha-
seolo Flos color Maluz, figura vero
Anemonis, folia octonæ lineæ rubræ
perreptant.

BANAICXITL.

A Napi radícula oriuntur quinæ. caules
rubri. rami omnes alati, Glycyrhizæ
modo ac figura. vltimum tamen fo-
liorum Thalictri, vel Adianto si-
mile.

CA-



ALIE PLANTÆ NOVÆ HISP.
CACACILI. CACALIC.



IOANNIS TERRENTII LYNCEI
CACACILI. CACAMILXOCHITL.

Radix longa, subnigra. caules foliati
Conyzæ instar, sine pediculo in sum-
mo botri Oxalidis foliorum viridis
diluti albicantis.

CACALIC.

Folium Cannabi simile, ferratum instar
Crithmi. nodi caulis magni. flos fla-
uus. calix squamosus.

CACAMILXOCHITL.

Vel *Thlacamilxochitl* *Quauhquiculle*.

Huic radix grumosa, Thoræ instar, co-
lore suberis. caulis bina habet folia
sibi opposita, Verbenaë similia. Flos
coccineus, in medio luteus, Rosæ si-
milis.



CA-

NARDI ANT. RECCHI. 359
CACAPATLI. CACATLAOLXOCHITL.



NOTATIONES.

CACHRVIAQVA.



CACAPATLI.

Pediculi longi ab radice orti, quibus in-
fident folia tria ferrata. Fragariae in-
star.

CACATLAOLXOCHITL.

Folia Castaneæ modo ferè ferrata. in-
cissione tamen non ita frequenti.

CACHRVIAQVA.

Ad Campanulas pertinere videtur. ca-
lix floris luteus, flos ipse ruber.

CA-

CAHVAZITZIQUI.

CAMOPALTIC.



IOANNIS TERRENTII LYNCEI

CAHVAZITZIQUI.

Radix longa, rubet instar Rhabarbar, flos longus carnei coloris, stamina lutea. herba est calida, & sicca, excessu tertio, & styptica.

CAMOPALTIC.

Caulis rubent. in summo flores multi simul, quiuis in superiore parte diuisi in quinque partes. inferiore tamen sui parte videtur esse vnicus.

CAMATOTONCAPATLI.

Caulis nigris maculis conspersus, folium cordis, cuius longitudo, & latitudo fere eadem, quatuor nempe digitorum.

CAMOL.



CHACHALACAME TL.

CAMOLXOCHITL.



NOTATIONES.

CENANAM DE QVAHVNA HOAC.

CHACHALACAME TL.

Est arbor baccifera: folia Pyro similia longa 4. lata 3. digitos: bacca sunt quales Ribium.

CAMOLXOCHITL.

Arbor leguminosa, folia in alas componit. Flos Cassia similis ex rubro flauus in 8. partes diuisus, siliqua. Piso similis.

CENANAM DE QVAHVNA HOAC.

Foliu Nerio simile, & foliolis ad maiorum alas erumpentibus, venisque in transuersum euntibus, Ora foliorum & vena media rubent. Bacca in vnam congesta virides sunt, vt in Solano.

Hh

CE-

