

## CONCLUSIONES

Los resultados obtenidos del presente estudio nos permitieron primeramente establecer que existe la presencia inmunológica de anticuerpos en el ganado bovino del Noreste de México (Coahuila, Nuevo León y Tamaulipas) producidos en contra de los antígenos pertenecientes al Protozoario *Neospora caninum*, agente causal de la denominada enfermedad de la Neosporosis. Cabe mencionar que fue encontrada tal evidencia serológica tanto en ganado lechero como de carne. Asimismo se reconfirmó que todavía existe la presencia serológica de otras enfermedades de importancia reproductiva y económica como la Brucellosis y la Infección por el Virus de la Leucosis Bovina en niveles por arriba de los esperados.

Las conclusiones sobre Neosporosis son las siguientes:

- ✓ La seroprevalencia en bovinos de la zona Noreste de México es de 36%.
- ✓ Los Estados más afectados fueron Coahuila con 45% de prevalencia, seguido por Nuevo León con 40% y el menos afectado fue Tamaulipas con 16%.
- ✓ La seroprevalencia por municipio estudiado varió desde 0 hasta 78%.
- ✓ Los bovinos destinados a la producción de leche resultaron más afectados con 39% que los destinados a la producción de carne con 10%.
- ✓ La raza Holstein-Friesian con 40% de prevalencia resultó la más afectada y las menos afectadas fueron las razas Charolais con 10% y Criollo con 0.

Las conclusiones con respecto a Leucosis son las siguientes:

- ✓ La seroprevalencia en bovinos lecheros de la zona Noreste de México es de 35%.
- ✓ El Estado más afectado fue Nuevo León con 52% y los menos afectados son Coahuila con 35% y Tamaulipas con 18%.
- ✓ La seroprevalencia por Municipio estudiado varió desde 0 hasta 100%.
- ✓ La raza Holstein-Friesian resultó ser la más afectada con 36% de prevalencia, mientras que los bovinos Criollos resultaron ser la más baja con 0.

Las conclusiones con respecto a Brucelosis son las siguientes:

- ✓ La seroprevalencia en bovinos lecheros de la zona Noreste de México es de 4.3%.
- ✓ El Estado más afectado fue Nuevo León con 6% y los menos afectados fueron Coahuila con 5% y Tamaulipas con 2%.
- ✓ La seroprevalencia por Municipio estudiado varió desde 0 hasta 40%.
- ✓ La raza Holstein-Friesian resultó ser la más afectada con 4.3% de prevalencia que los bovinos Criollos donde la prevalencia fue de 0.

Tales datos nos indican que existen fallas en los programas de manejo para el ganado, tanto de leche como para el de carne, por lo que hay que reforzar los conocimientos y esfuerzos en materia zoosanitaria para los veterinarios encargados de dichas instalaciones. Esto se pudiese lograr con las campañas de control y erradicación de las enfermedades en las diversas ganaderías en las

zonas afectadas para tener mejores condiciones ambientales, reconocer los factores de riesgo y alcanzar más altos niveles de producción.

## REFERENCIAS

- Abramova, E.N., Kondratev, V.S. and Stinstkii, I.A. 1979. The biochemistry of leucosis in cattle. Vet. Bulletin. 44: 689-704.
- Almansa, J.E., Mariño, O.C., Martínez, S.T., Rueda, O.C. 1996. Comparación preliminar entre dos métodos de producción de antígeno gp51 del virus de LEB. Rev. Téc. Científica. 7: 25-30.
- Altaner, C.1993. Envelope glycoprotein gp 51 of bovine leukemia virus is differently glycosilated in cells of various species and organ origin. Vet. Immunol. and Immunopathol. 36 : 163-177.
- Alton, G.G., Jones, L.M., Angus, R.D. and Verger, J.M. 1988. Techniques for the brucellosis laboratory. INRA. Paris. 25: 31-45.
- Anderson, M.L., Blanchard, P.C., Barr, B.C., Dubey, J.P., Hoffman, R.L. and Conrad, P.A. 1991. *Neospora*-like protozoan infection as a major cause of abortion in California dairy cattle. J. Am. Vet Med. Assoc. 198: 241-244.
- Ariza, J., Pellicer, T., Pallares, R. and Gudiol, F. 1992. Specific antibody profile in human brucellosis. Clin. Infect. Dis. 14: 131-140.
- Ávalos, R.R. 1995. "Detección serológica y molecular del virus de la leucosis bovina (VLB) en el Noreste de México". Tesis FCB/UANL, Monterrey, N. L.
- Barber, J.S., Van Ham, K. and Polis, I. 1997. Seroprevalence of antibodies to *Neospora caninum* in Belgian dogs. J. Small Anim. Pract. 38: 15-16.
- Barr, B.C., Anderson, M.L., Blanchard, P.C., Daft, B.M., Kinde, H. And Conrad, P.A. 1990. Bovine fetal encephalitis and myocarditis associated with protozoal infections. Vet. Pathol. 27: 354-361.

Barr, B.C., Anderson, M.L., Dubey, J.P. and Conrad, P.A. 1991. *Neospora*-like protozoal infections associated with bovine abortions. Vet. Pathol. 28: 110-116.

Barr, B.C., Anderson, M.L., Sverlow, K.W. and Conrad, P.A. 1995. Diagnosis of bovine fetal *Neospora* infection with an indirect fluorescent antibody test. Vet. Rec. 137: 611-613.

Barr, B.C., Anderson, M.L., Woods, L.W., Dubey, J.P. and Conrad, P.A. 1992. *Neospora*-like protozoal infections associated with abortion in goats. J. Vet. Diagn. Invest. 4: 365-367.

Barr, B.C., Conrad, P.A., Sverlow, K.W., Tarantal, A.F. and Hendrickx, A.G. 1994a. Experimental fetal and transplacental *Neospora* infection in the nonhuman primate. Lab. Invest. 71: 236-242.

Barr, B.C., Dubey, J.P., Lindsay, D.S., Reinolds, J.P. and Wells, S.J. 1998. Neosporosis: its prevalence and economic impact. Comp. Cont. Educ. Pract. Vet. Suppl. 20: 1-16.

Barr, B.C., Rowe, J.D., Sverlow, K.W., BonDurant, R.H., Ardans, A.A., Oliver, M.N. and Conrad, P.A. 1994b. Experimental reproduction of bovine fetal *Neospora* infection and death with a bovine *Neospora* isolate. J. Vet. Diagn. Invest. 6: 207-215.

Baszler, T.V., Knowles, D.P., Dubey, J.P., Gay, J.M., Mathison, B.A. and McElwain, T.F. 1996. Serological diagnosis of bovine neosporosis by *Neospora caninum* monoclonal antibody-based competitive inhibition ELISA. J. Clin. Microbiol. 34: 1423-1428.

Beal, V.C. 1984. Current estimated brucellosis losses. Veterinary Services. Animal and Plant Health Inspection Services. U.S. Department of Agriculture. Hyattsville, Md.

Bendixen, H.J. 1959. Studies on leukosis in cattle. Control of leukosis herds using hematological examination. Word .Vet. Med. 11: 733-738.

Bjerkas, I. and Dubey, J.P. 1991. Evidence that *Neospora caninum* is identical to the Toxoplasma-like parasite of Norwegian dogs. Acta Vet. Scand. 32: 407-410.

Bjerkas, I., Mohn, S.F. and Presthus, J. 1984. Unidentified cyst-forming sporozoan causing encephalomyelitis and myositis in dogs. Z. Parasitenkd. 70: 271-274.

Bjerkas, I. and Presthus, J. 1988. Immuno-histochemical and ultrastructural characteristics of a cyst-forming sporozoan associated with encephalomyelitis and myositis in dogs. Acta Pathol. Microbiol. Immunol. Scand. 96: 445-454.

Bjerkas, I. and Presthus, J. 1989. The neuropathology in toxoplasmosis-like infection caused by a newly recognized cyst-forming sporozoan in dogs. Acta Pathol. Microbiol. Immunol. Scand. 97: 459-468.

Bjorkman, C., Johnsson, O., Stenlund, S., Holmdahl, J. and Uggla, A. 1996. *Neospora* species infection in a herd of dairy cattle. J. Am. Vet. Med. Assoc. 208: 1441-1444.

Bjorkman, C., Lundén, A., Holmdahl, O.J.M., Barber, J., Trees, A.J. and Uggla, A. 1994. *Neospora caninum* in dogs: detection of antibodies by ELISA using an iscom antigen. Parasite. Immunol. 16: 643-648.

Bjorkman, C. and Uggla, A. 1999. Serological diagnosis of *Neospora caninum* infection. J. Parasitol. 29: 1497-1507.

Blood, D. y Rodostits, O. 1992. Leucosis viral bovina. En : Medicina Veterinaria. Edit. Interam. Mc Graw-Hill. México. 2: 878-886.

Bothwell, P.W. 1962. Economic losses of bovine brucellosis. Vet. Rec. 74: 1091.

Boulton, J.G., Gill, P.A., Cook, R.W., Fraser, G.C., Harper, P.A.W. and Dubey, J.P. 1995. Bovine *Neospora* abortion in north-eastern New South Wales. Aust. Vet. J. 72: 119-120.

Burny, A.C., C. Bruck, Y. Cleuter, D. Couez, J. Deschamps, J. Ghysdael, D. Gregoire, R. Kettmann, M. Mammerickx, G. Marbaix, D. Portetelle, L. Willems. (1985). Bovine leukemia virus, a versatile agent with various pathogens effects in various animals species. Cancer Research (suppl.) Cancer Research 45:4578s-4582s.

Burridge, M.J., Puhr, M.D. and Hennemann, J.M. 1981. Prevalence of bovine leukemia virus infection in Florida. J. Am. Vet. Med. Assoc. 179: 704-707.

Buxton, D., Caldow, G.L., Maley, S.W., Marks, J. and Innes E.A. 1997. Neosporosis and bovine abortion in Scotland. Vet. Rec. 141: 649-651.

Buzala, E. and Deren, W. 2003. Comparison of PLA with AGID and ELISA results in serology diagnosis of bovine leukosis. Pol. J. Vet. Sci. 6: 9-11.

- Callis, J., Bachrach, H., Bittle, J., Dalrymple, J., Gamble, R., Glosser, J., Murphy, F., Thiermann, A. and Thompson, S. 1989. Biotechnology and its public health implications in zoonotic diseases. *Biotechnology for livestock production.* New York, Plenum Press. 25: 377-400.
- Calzada, C.P., Morales, S.E., Quiroz, R.G., Salmerón, S.F., García, O.C., Hernández, B.J. 2002. Valores hematológicos en vacas de raza Holstein Friesian seropositivas a *Neospora caninum* de la cuenca lechera de Tizayuca, Hidalgo, México. Vet. Mex. 33:119-124.
- Cerqueira, L.R., Modena, C.M., Moreira, E.C. and Abreu, J.J. 1984. Evolucao clínica de Leucose Enzoótica Bovina. Arq. Bras. Med. Vet. Zoot. 36: 47-57.
- Chasey, D., Wibberley, G., Markson, M. and Roberts, D.H. 1978. Demostration of enzootic bovine leucosis virus in Britain. Vet. Rec. 103: 449-450.
- Chi, J., VanLeeuwen, J.A., Weersink, A. and Keefe, G.P. 2002. Direct production losses and treatment costs from bovine viral diarrhoea virus, bovine leukosis virus, *Mycobacterium avium* subspecies paratuberculosis, and *Neospora caninum*. Prev. Vet. Med. 55: 137-153.
- Cole, R.A., Lindsay, D.S., Blagburn, B.L. and Dubey, J.P. 1995a. Vertical transmission of *Neospora caninum* in mice. J. Parasitol. 81: 730-732.
- Cole, R.A., Lindsay, D.S., Blagburn, B.L., Sorjonen, D.C. and Dubey, J.P. 1995b. Vertical transmission of *Neospora caninum* in dogs. J. Parasitol. 81: 208-211.

- Cole, R.A., Lindsay, D.S., Dubey, J.P., Toivio-Kinnucan, M.A. and Blagburn, B.L. 1994. Characterization of a murine monoclonal antibody generated against *Neospora caninum* by western blot analysis and immunoelectron microscopy. Am. J. Vet. Res. 55: 1717-1722.
- Conrad, P.A., Sverlow, K., Anderson, M., Rowe, J., BonDurant, R., Tuter, G., Breitmeyer, R., Palmer, C., Thurmond, M., Ardans, A., Dubey, J.P., Duhamel, G. and Barr, B. 1993. Detection of serum antibody responses in cattle with natural or experimental *Neospora* infections. J. Vet. Diagn. Invest. 5: 572-578.
- Crawford, R.P., Williams, J.D., Huber, J.D. and Childers, A.B. 1973. Biotypes of *Brucella abortus* and their value in epidemiological studies of infected cattle herds. J. Am. Vet. Med. Assoc. 175: 121.
- Cuddon, P., Lin, D.-S., Bowman, D.D., Lindsay, D.S., Miller, T.K., Duncan, I.D., deLahunta, A., Cummings, J., Suter, M., Cooper, B., King, J.M. and Dubey, J.P. 1992. *Neospora caninum* infection in English Springer Spaniel Littermates: Diagnostic evaluation and organism isolation. J. Vet. Intern. Med. 6: 325-332.
- Dannatt, L. 1997. *Neospora caninum* antibody levels in an endemically-infected dairy herd. Cattle pract. 5: 335-337.
- Davison, H.C., Otter, A. and Trees, A.J. 1999. Estimation of vertical and horizontal transmission parameters of *Neospora caninum* infections in dairy cattle. Int. J. Parasitol. 29 : 1683 –1689.

Delgado, G.R., Quintero, C.J., De Luna, A. A. 1995. Estudio patológico microbiológico y serológico del aborto en bovinos Holstein de la Comarca Lagunera. Memorias del XIX Congreso Nacional de Buiatría; agosto 24-26; Torreón, Coahuila, México. Asociación Mexicana de Médicos Veterinarios Especialistas en Bovinos, AC, 1995: 74-78.

Dreier, K.J., Warren Stewarter, L., Kerlin, R.L., Ritter, D.M. and Brake, D.A. 1999. Phenotypic characterization of a *Neospora caninum* temperature-sensitive strain in normal and immunodeficient mice. 29: 1627-1634.

Deshayes, L., Levy, D., Parodi, A.L. and Levy, J.P. 1980. Spontaneus immune response of bovine leukemia virus infected cattle against five different viral proteins. Int. J.Cancer. 25: 503-508.

Dubey, J.P. 1992. A review of *Neospora caninum* and Neospora-like infections in animals. J. Protozool. Res. 2:40-52.

Dubey, J.P. 1999. Neosporosis a newly recognized protozoan disease similar to Toxoplasmosis. The Infectious Disease Review. 2:129-130.

Dubey, J.P. 2003. Review of *Neospora caninum* and neosporosis in animals. Korean J. Parasitol. 41: 1-16.

Dubey, J.P. 1993. *Toxoplasma, Neospora, Sarcocystis*, and other tissue cyst-forming coccidia of humans and animals. In: Kreier J.P. (Editor). Parasitic Protozoa. 6: 1-158.

Dubey, J.P., Carpenter, J.L., Speer, C.A., Topper, M.J., Uggla, A. 1988a. Newly recognized fatal protozoan disease of dogs. J. Am. Vet. Med. Assoc. 192:1269-85.

Dubey, J.P. and de Lahunta, A. 1993. Neosporosis associated congenital limb deformities in a calf. Appl. Parasitol. 34: 229-233.

Dubey, J.P., Hartley, W.J. and Lindsay, D.S. 1990a. Congenital *Neospora caninum* infection in a calf with spinal cord anomaly. J. Am. Vet. Med. Assoc. 197: 1043-1044.

Dubey, J.P., Hartley, W.J., Lindsay, D.S. and Topper, M.J. 1990b. Fatal congenital *Neospora caninum* infection in a lamb. J. Parasitol. 76: 127-130.

Dubey, J.P., Hattel, A.L., Lindsay, D.S. and Topper, M.J. 1988b. Neonatal *Neospora caninum* infection in dogs: Isolation of the causative agent and experimental transmission. J. Am. Vet. Med. Assoc. 193: 1259-1263.

Dubey, J.P. and Lindsay, D.S. 1996. A review of *Neospora caninum* and neosporosis. Vet. Parasitol. 67:1-59.

Dubey, J.P. and Lindsay, D.S. 1990a. Neosporosis in dogs. Vet. Parasitol. 36: 147-151.

Dubey, J.P. and Lindsay, D.S. 1990b. *Neospora caninum* induced abortion in sheep. J. Vet. Diagn. Invest. 2: 230-233.

Dubey, J.P. and Lindsay, D.S. 1989a. Transplacental *Neospora caninum* infection in cats. J. Parasitol. 75: 765-771.

Dubey, J.P. and Lindsay, D.S. 1989b. Transplacental *Neospora caninum* infection in dogs. Am. J. Vet. Res. 50: 1578-1579.

Dubey, J.P. and Lindsay, D.S. 1989c. Fatal *Neospora caninum* infection in Kittens. J. Parasitol. 75: 148-151.

Dubey, J.P., Lindsay, D.S., Adams, D.S., Gay, J.M., Baszler, T.V., Blagburn, B.L. and Thulliez, P. 1996. Serologic responses of cattle and other animals infected with *Neospora caninum*. Am. J. Vet. Res. 57: 329-336.

Dubey, J.P., Lindsay, D.S., Anderson, M.L., Davis, S.W. and Shen, S.K., 1992. Induced transplacental transmission of *Neospora caninum* in cattle. J. Am. Vet. Med. Assoc. 201: 709-713.

Dubey, J.P., Lindsay, D.S., Hill, D., Romand, S., Thulliez, P., Kwok, O.C.H., Silva, J.C.R., Oliveira-Camargos M.C. and Gennari, S.M. 2002. Prevalence of antibodies to *Neospora caninum* and *Sarcocystis neurona* in sera of domestic cats from Brazil. J. Parasitol. 88: 1251-1252.

Dubey, J.P., Metzger, F.L.J., Hattel, A.L., Lindsay, D.S. and Fritz, D.L. 1995. Canine cutaneous neosporosis: clinical improvement with clindamycin. Vet. Dermatol. 6: 37-43.

Dubey, J.P., Miller, S., Lindsay, D.S. and Topper, M.J. 1990c. *Neospora caninum*-associated miocarditis and encephalitis in an aborted calf. J. Vet. Diagn. Invest. 2: 66-69.

Dubey, J.P. and Porterfield, M.L. 1990. *Neospora caninum* (Apicomplexa) in an aborted equine fetus. J. Parasitol. 76: 732-734.

Espada, R., Flogio, A., Gurría, F., López, G., López, J., Meixueiro, H., Pérez, J., Yáñez, V., Hernández, O., Beymer, D., Riemman, H. 1986. Prevalencia de anticuerpos contra las enfermedades infecciosas más comunes del ganado bovino de Baja California. Vet. Mex. 17: 23-29.

Esteban, E.N., Thorn, R.M. and Ferrer, J.F. 1985. Characterization of the blood lymphocyte population in cattle infected with the bovine leukemia virus. Cancer Research. 45: 3225-3230.

Fekete, A., Bantle, J.A. and Halling, S.M. 1992. Detection of Brucella by polymerase chain reaction in bovine fetal and maternal tissues. J. Vet. Diagn. Invest. 4: 79-83.

Ferrer, J.F., Marshak, R.R., Abt, D.A. and Kenyon, S. 1978. Persistent lymphocytosis in cattle: its cause, nature and relation to lymphosarcoma. Ann. Rech. Vet. 9: 851-857.

Gonzalez, L., Buxton, D., Atxaerandia, R., Aduriz, G., Maley, S., Marco, J.C. and Cuervo, L.A. 1999. Bovine abortion associated with *Neospora caninum* in northern Spain. Vet. Rec. 144: 145-150.

Gottstein, B., Henrich, B., Wyss, R., Thur, B., Busato, A., Stark, K.D.C. and Muller, N. 1998. Molecular and immunodiagnostic investigations on bovine neosporosis in Switzerland. Int. J. Parasitol. 28: 679-691.

Goudswaard, J., Gilmourt, N.J.L., Dijkstra, R.G. and Beck, J.J.V. 1976. Diagnosis of johne's disease in cattle; a comparison of five serological tests under field conditions. Vet. Rec. 98: 461-462.

Guarino, A., Fusco, G., Savini, G., Di Francesco, G. and Cringoli, G. 2000. Neosporosis in water buffalo (*Bubalus bubalis*) in Southern Italy. Vet. Parasitol. 91: 15-21.

Gurturk, K., Boynukara, B., Lihan, Z., Hakki Ekin, I. and Gulhan, T. 1999. Comparison of the dot-immunobinding assay with the serum agglutination test, the rose Bengal plate test and the milk ring test for the detection of

Brucella antibodies in bovine sera and milk. Zentralbl Veterinarmed B. 46: 279-285.

Hagan, Bruner. 1983. Enfermedades infecciosas de los animales domésticos. 4ta edición. Editorial la Prensa Mexicana. p.106-129.

Harrington, R. and Brown, G.M. 1976. Laboratory summary of *Brucella* isolations and typing. Am. J. Vet. Res. 37: 1241.

Hay, W.H., Shell, L.G., Lindsay, D.S. and Dubey, J.P. 1990. Diagnosis and treatment of *Neospora caninum* infection in dogs. J. Am. Vet. Med. Assoc. 197: 87-89.

Hemphill, A., Gottstein, B. and Kaufmann, H. 1996. adhesion and invasion of bovine endothelial cells by *Neospora caninum*. Parasitol. 112: 183-197.

Hietala, S. K. and Thurmond, M.C. 1999. Postnatal *Neospora caninum* transmission and transient serologic responses in two dairies. International Journal of Parasitology. 29:1669-1676.

Ho, M.S.Y., Barr, B.C., Tarantal, A.F., Lai, L.T.Y., Hendrickx, A.G., Marsh, A.E., Sverlow, K.W., Packham, A.E. and Conrad, P.A. 1997. Detection of *Neospora* from tissues of experimentally infected Rhesus Macaques by PCR and specific DNA probe hybridization. Am. Soc. Microbiol. 35: 1740-1745.

Jacobs, R.M. 1983. Bovine lymphoma. In Comparative Pathobiology of Viral Diseases. 2: 21-51.

Jardine, J.E. 1996. The ultrastructure of bradyzoites and tissue cysts of *Neospora caninum* in dogs: absence of distinguishing morphological features between parasites of canine and bovine origin. Vet. Parasitol. 62: 231-240.

Jensen, L., Jensen, T.K., Lind, P., Henriksen, S.A., Uggla, A. and Bille-Hansen, V. 1998. Experimental porcine neosporosis. APMIS. 106: 475-482.

Johnson, R., Gibson, C.D. and Kaneene, J.B. 1985. Bovine leukemia virus: a herd-based control strategy. Prev. Vet. Med. 3: 339-349.

Johnson, R. and Kaneene, J.B. 1991a. Bovine leukemia virus. Part I. Descriptive epidemiology, clinical manifestations and diagnostic test. Comp. Cont Ed. 13: 315-327.

Johnson, R. and Kaneene, J.B. 1991b. Bovine leukemia virus. Part II. Risk factor of transmission. Comp. Cont Ed. 13: 681-691.

Kaaden, O. and Stephenson, J.R. 1978. Report of the second session detection of BLV infection: serological methods. Ann. Res. Vet. 9: 604-605.

Kasari, T.R., Barling, K. and McGrann, J.M. 1999. Estimated production and economical losses from *Neospora caninum* infection in Texas beef-herds. The Bovine Practitioners 33:113-120.

Kuby, J. 1992. Antigen-antibody interactions. Immunol. 1st ed. W.H. Freem. Comp. NY, 6: 126-135.

Lassauzet, M.G., Thurmond, M.C., Johnson, W.O., Stevens, F. and Picanso, J.P. 1991. Factor associated with transmission of bovine leukemia virus by contact in cows on a California dairy. Am. J. Epidemiol. 133: 164-176.

- Lindsay, D.S., Blagburn, B.L. and Dubey, J.P. 1992. Factors affecting the survival of *Neospora caninum* bradizoites in murine tissues. *J. Parasitol.* 78: 70-72.
- Lindsay, D.S. and Dubey, J.P. 1989a. In vitro development of *Neospora caninum* (Protozoa: Apicomplexa) from dogs. *J. Parasitol.* 75: 163-165.
- Lindsay, D.S. and Dubey, J.P. 1989. Immunohistochemical diagnosis of *Neospora caninum* in tissue sections. *Am. J. Vet. Res.* 50:1981-1983.
- Lindsay, D.S. and Dubey, J.P. 1990. *Neospora caninum* (Protozoa: Apicomplexa) infections in rats. *Can. J. Zool.* 68: 1595-1599.
- Lindsay, D.S., Dubey, J.P. and Blagburn, B.L. 1991. Characterization of a *Neospora caninum* (Protozoa. Apicomplexa) isolate (NC-3) in mice. *J. Alabama Acad. Sci.* 62: 1-8.
- Lindsay, D.S., Dubey, J.P. and McAllister, M.M. 1999. *Neospora caninum* and the potential for parasite transmission. *Small Anim. Exot.* 21: 317-321.
- Lindsay, D.S., Kelly, E.J., McKown, R., Stein, F.J., Plozer, J., Herman, J., Blagburn, B.L. and Dubey, J.P. 1996a. Prevalence of *Neospora caninum* and *Toxoplasma gondii* antibodies in coyotes (*Canis latrans*) and experimental infections of coyotes with *Neospora caninum*. *J. Parasitol.* 82: 657-659.
- Lindsay, D.S., Little, S.E. and Davidson, W.R. 2002. Prevalence of antibodies to *Neospora caninum* in white-tailed deer, *Odocoileus virginianus*, from the southeastern United States. *J. Parasitol.* 88: 415-417.

- Lindsay, D.S., Rippey, N.S., Powe, T.A., Sartin, E.A., Dubey, J.P. and Blagburn, B.L. 1995. Abortions, fetal death, and stillbirths in pregnant pygmy goats inoculated with tachizoites of *Neospora caninum*. Am. J. Vet. Res. 56: 1176-1180.
- Lindsay, D.S., Speer, C.A., Toivio-Kinnucan, M.A., Dubey, J.P. and Blagburn, B.L. 1993. Use of infected cultured cells to compare ultrastructural features of *Neospora caninum* from dogs and *Toxoplasma gondii*. Am. J. Vet. Res. 54: 103-106.
- Lindsay, D.S., Spencer, J. and Rupprecht, C. 2001. Prevalence of agglutinating antibodies to *Neospora caninum* in raccoons, *Procyon lotor*. J. Parasitol. 87: 1197-1198.
- Lindsay, D.S., Steinberg, H., Dubielzig, R.R., Semrad, S.D., Konkle, D.M., Miller, P.E. and Blagburn, B.L. 1996b. Central nervous system neosporosis in a foal. J. Vet. Diagn. Invest. 85: 658-660.
- Locatelli-Dittrich, R., Soccol, V.T. and Richartz, R.R.T.B. 2001. Serological diagnosis of neosporosis in a herd of dairy cattle in southern Brazil. J. Parasitol. 87: 1493-1494.
- Luna-Martinez, J.E. and Mejia-Teran, C. 2002. Brucellosis in Mexico: current status and trends. Vet. Microbiol. 90: 19-30.
- Mainar-Jaime, R.C., Thurmond, M.C., Berzal-Herranz B., Hietala, S.K. 1999. Seroprevalence of *Neospora caninum* and abortion in dairy cows in northern Spain. Vet. Rec. 145:72-75.

- Mamoun, R.Z., Astier, T., Guilleman, B. and Duplan, J.F. 1983. Bovine lymphosarcoma: Processing of bovine leukemia virus-coded proteins. J. Gen. Virol. 64: 2791-2795.
- Mariño, O.C., Bravo, M.M., Fajardo, E. 1984. La técnica de ELISA en el diagnóstico serológico de la Leucosis bovina en Colombia. Rev. ICA. 19: 315-323.
- Marsh, A.E., Howe, D.K., Wang, G., Barr, B.C., Cannon, N. and Conrad, P.A. 1999. Differentiation of *Neospora hughesi* from *Neospora caninum* based on their immunodominant surface antigen, SAG1 and SRS2. Int. J. Parasitol. 29: 1575-1582.
- Mayhew, I.G., Smith, K.C., Dubey, J.P., Gatward, L.K. and McGlenon, N.J. 1991. Treatment of encephalomyelitis due to *Neospora caninum* in a litter of puppies. J. Small Anim. Pract. 32: 609-612.
- McAllister M.M., Dubey J.P., Lindsay D.S., Jolley W.R., Wills R.A. McGuire A.M. 1998. Dogs are definitive hosts of *Neospora caninum*. Int. J. Parasitol. 28:1473-1478.
- McAllister M.M., Parmley, S.F., Weiss, L.M., Welch, V.J. and McGuire, A.M. 1996. An immunohistochemical method for detecting bradyzoite antigen. J. Parasitol. 82: 354-355.
- McGuire, A.M., McAllister, M.M., Wills, R.A. and Tranas, J.D. 1999. Experimental inoculation of domestic pigeons (*Columba livia*) and Zebra finches (*Poephila guttata*) with *Neospora caninum* tachizoites. Int. J. Parasitol. 29: 1525-1529.

McNamee, P.T., Trees, A.J., Guy, F., Moffett, D. and Kilpatrick, D. 1996. Diagnosis and prevalence of neosporosis in cattle in northern Ireland. Vet. Rec. 138: 419-420.

Meas, S., Ruas, J., Farias, N.A., Usui, T., Teraoka, Y., Mulenga, A., Chang, K.S., Masuda, A., Madruga, C.R., Ohashi, K. and Onuma, M. 2002. Seroprevalence and Molecular evidence for the presence of bovine immunodeficiency virus in brazilian cattle. Jpn. J. Vet. Res. 50: 145.

Meas, S., Yilmaz, Z., Usui, T., Torun, S., Yesilbag, K., Ohashi, K. and Onuma, M. 2003. Evidence of bovine immunodeficiency virus in cattle in Turkey. Jpn. J. Vet. Res. 51: 3-8.

Milner, A.R., Mack, W.N.M., Coates, K.J., Hill, J., Gill, I. and Sheldrick, P. 1990. The sensitivity and specificity of a modified ELISA for the diagnosis of Johne's disease for a field trial in cattle. Vet. Microbiol. 25: 193-198.

Mirsky, M.L., Olmstead, C. and Lewin, H. 1996. Prevalence of proviral bovine leukemia virus in peripheral blood mononuclear cells at two subclinical stages of infection. J. Virol. 70: 2178-2183.

Moen, A.R., Wouda, W., Mul, M.F., Graat, E.A.M. and van Werven, T. 1998. Increased risk of abortion following *Neospora caninum* abortion outbreaks: a retrospective and prospective cohort study in four dairy herds. Theriogenology. 49:1301-1309.

Monroy, B.J., Trigo, T.F., Larios, G.F., Fajardo, M.R., Marquez. 1985. Estudio seroepidemiológico de Leucosis bovina en México. Memorias de la reunión de investigación pecuaria en México. Mex. D.F. p. 84.

- Morales, S.E., Ramirez, L.J., Trigo, T.F., Ibarra, V.F., Puente, C.E., Santacruz, M. 1997. Descripción de un caso de aborto Bovino asociado a infección por *Neospora sp* en México. Vet. Méx. 28:353-357.
- Morales, S.E., Trigo, T.F., Ibarra, V.F., Puente, C.E., SantaCruz, M. 2001. Seroprevalence study of bovine neosporosis in México. J. Vet. Diagn. Invest. 13:413-415.
- Moreno, E. 2002. Brucellosis in Central America. Vet. Microbiol. 90: 31-38.
- Nicoletti, P. 1980. The epidemiology of bovine brucellosis. Adv. Vet. Sci. Comp. Med. 24: 69-98.
- Noda, J., Pérez, M., Marrera, M., Abeledo, M.A. 1992. The use of ELISA and Nucleid acid hybridization test in research and diagnosis of bovine leukosis virus. JAVMA. 199: 85-90.
- Nuotio, L., Rusanen, H., Sihuonen, L. and Neuvonen, E. 2003. Eradication of LEB from Finland. J. Vet. Med. Sci. 65: 287-289.
- Olds, R.J. 1982. Atlas de microbiología. Editorial Wolfe, científico médica. Primera edición, España. Pp. 72-73.
- Ollis, G. 1996. Enzootic bovine leukosis in dairy cattle. J. Virol. Meth. 41: 101-112.
- Olson, C. and Miller, J.M. 1987. History and Terminology of enzootic bovine leukosis and bovine leukemia virus. JAVMA. 155: 3-11.
- Omer, M.K., Assefaw, T., Skjerve, E., Tekleghiorghis, T. and Woldehiwet, Z. 2002. Prevalence of antibodies to *Brucella spp.* And risk factors related to high – risk occupational groups in Eritrea. Epidemiol. Infect. 129: 85-91.

Osawa, T., Wastling, J., Maley, S., Buxton, D. and Innes, E.A. 1998. A multiple antigen ELISA to detect *Neospora*-specific antibodies in bovine sera, bovine fetal fluids, ovine and caprine sera. Vet. Parasitol. 79: 19-34.

Otter, A., Griffiths, I.B. and Jefrey, M. 1993. Bovine *Neospora caninum* abortion in the UK. Vet. Rec. 133: 375.

Paré J., Hietala, S.K. and Thurmond, M.C. 1995. An enzyme-linked immunosorbent assay (ELISA) for serological diagnosis of *Neospora* sp. infection in cattle. J. Vet. Diagn. Invest. 7: 352-359.

Paré J., Thurmond M.C. and Hietala S.K. 1996. Congenital *Neospora caninum* infection in dairy cattle and associated calfhood mortality. Can. J. Vet. Res. 60:133-139.

Paré J., Thurmond M.C. and Hietala S.K. 1994. Congenital *Neospora* infection in dairy cattle. Vet. Rec. 134: 531-532.

Payne, S. and Ellis, J. 1996. Detection of *Neospora caninum* DNA by the polymerase chain reaction. Int. J. Parasitol. 26: 347-351.

Pelzer, K. and Sprecher, D. 1993. Controlling BLV infection on dairy operations. Vet. Med. 25: 275-281.

Perry, M. and Bundle, D.R. 1990. Antigenic relationships of the lipopolysaccharides of *Escherichia hermanii* strains with those of *Escherichia coli* 0157:H7, *Brucella melitensis* and *Brucella abortus*. Inf. Immun. 58: 1391-1395.

- Peters, M., Wohlsein, P., Knieriem, A. and Schares, G. 2001. *Neospora caninum* infection associated with stillbirths in captive antelopes (*Tragelaphus imberbis*). Vet. Parasitol. 97: 153-157.
- Quinn, P.J., Carter, M.E., Markey, B. and Carter, G.R. 1994. Clinical veterinary microbiology. Editorial Wolfe. Primera edición, España. Pp. 262.
- Rebhun, W.C. 1982. Orbital lymphosarcoma in cattle. J. Am. Vet. Med. Assoc. 180: 149-152.
- Reed, V.I. 1981. Enzootic Bovine Leukosis. Can. Vet. J. 22: 95-102.
- Ressang, A.A., Gielkens, A.L.J., Quak, S., Mastenbroek, N., Tuppert, C. and De Castro, A. 1978. Enzyme linked immunosorbent assay for the detection of antibodies to bovine leukosis virus. Ann. Rech. Vet. 9: 663-666.
- Rhodes, J.K., Pelzer, K.D. and Johnson, Y.J. 2003. Economic implications of bovine leukemia virus infection in mid-Atlantic dairy herds. J. Am. Vet. Med. Assoc. 223: 346-352.
- Ruiz-Castenada, M. 1947. Studies on the pathogenesis of brucellosis. Proc. Soc. Exp. Biol. Med. 64: 241-244.
- Russel, A., Runnels, W.S. and Monlux, A.W. 1978. Principios de patología veterinaria. CECSA. p. 32.
- Schares, G., Peters, M., Wurm, R., Tackmann, K., Henning, K. and Conraths, F.J. 1997. *Neospora caninum* causes abortions in a cattle herd in North Rhine Westphalia. Deu. Tier. Wochensch. 104: 208-212.

Simpson, V.R., Monies, R.J. and Riley, P. 1997. Foxes and neosporosis. Vet. Rec. 141: 503.

Speer, C.A. and Dubey, J.P. 1989. Ultrastructural of tachizoites, bradizoites and tissue cyst of *Neospora caninum*. J. Protozool. 36: 458-463.

Spencer, J.A., Witherow, A.K. and Blagburn, B.L. 2000. A random amplified polymorphic DNA polymerase chain reaction technique that differentiates between *Neospora* species. Int. J. Parasitol. 30: 1366-1368.

Suzan, V.M., Onuma, M., Aguilar, R.E. and Murakawi, Y. 1983. Prevalence of bovine Herpesvirus-1, Parainfluenza-3, bovine Rotavirus, bovine viral diarrhea, bovine Adenovirus-7, bovine leukemia virus and bluetongue virus antibodies in cattle in México. Jpn. J. Vet. Res. 31: 125-132.

Stenlund, S., Kindahl, H., Magnusson, U., Uggla, A. and Bjorkman, C. 1999. Serum antibody profile and reproductive performance during two consecutive pregnancies of cows naturally infected with *Neospora caninum*. Vet. Parasitol. 85: 227-234.

Straub, O.C. and Olson, C. 1978. Report of the fourth session (Epidemiology of BLV). Third international symposium on bovine leukosis. Ann. Rech. Vet. 9: 605.

Teich, N. 1982. Taxonomy of retroviruses. In: Weiss R, Teich N, Varmus H, Coffin J. Eds .RNA Tumor Virus. Cold spring Harbor. NY: Cold Spring Harbor Laboratory: 25-28.

Thornton, R.N., Thompson, E.J. and Dubey J.P. 1991. *Neospora* abortion in New Zealand cattle. N. Z. Vet. J. 39:129-133.

Thurmond, M. and Hietala, S. 1997. Effect of congenitally acquired *Neospora caninum* infection on risk of abortion and subsequent abortions in dairy cattle. Am. J. Vet. Res. 58: 1381-1385.

Thurmond, M. and Hietala, S. 1995. Strategies to control *Neospora* infection in cattle. Bovine Pract. 29: 60-63.

Thurmond, M., Hietala, S., Blanchard, P. and Debey, S. 1995. Congenital transmission of *Neospora caninum* in Herds experiencing endemic or epidemic abortion. Proc. 38th Am. Meet. Am. Assoc. Vet. Lab. Diagn. Sparks. Nevada, p. 97.

Thurmond, M.C., Maden, C.B. and Carter, R.L. 1985. Cull rates of dairy cattle with antibodies to bovine leukemia virus. Canc. Res. 45:1987-1989.

Thoen, C.O. and Enright, F. 1986. *Brucella*. In, C.L. Gyles and C.O. Thoen (eds.) Pathogenesis of bacterial infections in animals. Ames. Iowa State University Press.

Trees, A.J., Guy, F., Low, J.C., Roberts, L., Buxton, D. and Dubey, J.P. 1994. Serological evidence implicating *Neospora* species as a cause of abortion in British cattle. Vet. Rec. 134: 405-407.

Uggla, A., Stenlund, S., Holmdahl, O.J.M., Jakubek, E.-B., Thebo, P., Kindahl, H. and Bjorkman, C. 1998. Oral *Neospora caninum* inoculation of neonatal calves. J. Parasitol. 28: 1467-1472.

Usui, T., Meas, S., Konnai, S., Ohashi, K. and Onuma, M. 2003. Seroprevalence of bovine immunodeficiency virus and bovine leukemia virus in dairy and beef cattle in hokkaido. J. Vet. Med. Sci. 65: 287-289.

Venturini, M.C. 1999. *Neospora caninum* infections in bovine foetuses and dairy cows with abortions in Argentina. Int. J. Parasitol. 29:1075-1708.

Voller, A., Bidwell, D.E. and Bartlett, A. 1982. ELISA techniques in virology in: New development in practical virology. p. 59-81.

Waldner, Ch.L., Janzen, E.D., Henderson, J. and Haines, D.M. Outbreak of abortion associated with *Neospora caninum* infection in a beef herd. Departments of Herd Medicine and Theriogenology and Microbiology, Western College of Veterinary Medicine, University of Saskatchewan.

Waldner, Ch.L., Janzen, E.D. and Ribble, C.S. 1998. Determination of the association between *Neospora caninum* infection and reproductive performance in beef herds. J. Am. Vet. Med. Assoc. 213: 685-690.

Williams, D.J.L. 1999. Evaluation of a commercial ELISA for detecting serum antibody to *Neospora caninum* in cattle. Vet. Rec. 145: 571-575.

Wouda, W., Bartels, C.J.M. and Moen, A.R. 1999a. Characteristics of *Neospora caninum*-associated abortion storms in dairy herds in the Netherlands (1995 to 1997). Theriogenology. 52: 233-245.

Wouda, W., Dijkstra, Th., Kramer, A.M.H., Van Maanen, C., and Brinkhof, J.M.A. 1999b. Seroepidemiological evidence for a relationship between *Neospora caninum* infections in dogs and cattle. Int. J. Parasitol. 29: 1677-1682.

Wouda, W., Dubey, J.P. and Jenkins, M.C. 1997. Serological diagnosis of bovine fetal neosporosis. J. Parasitol. 83: 545-547.

Wouda, W., Moen, A.R., Visser, I.J.R. and van Knapen, F. 1996. Bovine fetal neosporosis: A comparison of epizootic and sporadic abortion cases and different age classes with regards to lesion severity and immunohistochemical identification in brain, heart and liver. J. Vet. Diagn. Invest. 121: 397-405.

Wouda, W., van den Ingh, T.S.G.A.M., van Knapen, F., Sluyter, F.J.H., Koeman, J.P. and Dubey, J.P. 1992. *Neospora abortus* bij het rund in Nederland. Tijdschr. Diergeneeskdl. 117: 599-602.

Yamage, M., Flechtner, O. and Gottstein, B. 1996. *Neospora caninum*: specific oligonucleotide primers for the detection of brain "cyst" DNA of experimentally infected nude mice by the polymerase chain reaction (PCR). J. Parasitol. 82: 272-279.

Young, E.J. and Corbel, M.J. (eds.). 1989. Brucellosis: Clinical and laboratory aspects. Boca Ratón, Florida, CRC Press, INC.